

**Interspecific *Nicotiana* Hybrids and Their Progeny**  
**CROSS-REFERENCE TO RELATED APPLICATIONS**

10-10-68

**[0002]** The present invention relates to novel interspecific *Nicotiana excelsior* x *N. benthamiana* hybrid seeds and plants and to a method of producing interspecific *Nicotiana* hybrids having enhanced properties for biomass and the production of recombinant proteins using a viral vector system.

**[0003]** The difference between *intra*- and *interspecific* hybrids can be visualized by the Mendelian behavior of intraspecific hybrids and the essentially non-Mendelian behavior of interspecific hybrids. This difference results from the complete chromosome homology in hybrids between plants of the same species (intraspecific) and the partial or nonchromosome homology that characterizes hybrids between plants of different species (interspecific). The intermediate or gray area between the extremes among interspecific hybrids is seen in the behavior of those involving closely related species. Such species will have the same chromosome number, and the extent of chromosome homologies may be very high. Hybrids of this kind produce seed when self-pollinated and show evidence of Mendelian patterns of segregation for some traits.

**[0004]** Cytological studies of meiosis in some  $F_1$  hybrids may show evidence of chromosome irregularities that reflect the chromosomal differences that mark the parents as different species. A hybrid between distantly related species may show reduced pairing between chromosomes of the different genomes. Meiosis in such hybrids may exhibit the typical chromosome behavior characteristic of monogenomic haploids. Most interspecific  $F_1$  hybrid combinations in the genus *Nicotiana* fall into this latter category. Their chromosome doubled counterparts are called amphidiploids, or

simply allopolyploids, when a parent of the hybrid may itself be a combination of different ancestral genomes. *N. tabacum* represents a typical example.

**[0005]** The relative difficulty of producing interspecific  $F_1$  hybrids increases in proportion to the decrease in taxonomic relations between their parents. Although some  $F_1$  hybrids are stable when converted to fertile polyploids, even hybrids between distantly related parents share some degree of chromosome homology. There are advantages to maintaining seed stocks of allopolyploids by self-pollination, particularly if the hybrid is difficult to obtain by conventional cross-pollination.

**[0006]** Interspecific hybridization and introgression in natural populations of plants and animals is a known source of genetic variation and adaptation. The term introgression, or introgressive transfer, of genetic traits is used in a narrower sense with short-term breeding projects. The goal of experimental interspecific hybridization is usually the introgressive transfer of a specific trait from one species (the nonrecurrent parent) into the genome of another (the recurrent parent). The trait must be expressed with reasonable phenotypic fidelity as a dominant or partial dominant in the  $F_1$  hybrid and all subsequent backcross generations, when nonrecurrent chromosomes are lost at random, otherwise the proposed interspecific transfer could not be made. Foreign genes from one species can become unstable when translocated into the germplasm of another.

**[0007]** Chromosome pairing between the foreign genomes in an interspecific hybrid may be extensive or minimal although the  $F_1$  hybrid is usually sterile. Fertility may be restored by treating germinating seed of the hybrid in 0.4 percent aqueous colchicine for about 4 hours to induce chromosome doubling. Immersion of the seeds in 0.1 percent colchicine for 24 hours may also be effective. Thereafter, the seed is rinsed in sterile distilled water and planted by sowing on the surface of pasteurized soil in glass or plastic preparation dishes. The seedlings later are transplanted to pots of soil in the greenhouse and mature plants that show good pollen development are self pollinated or backcrossed to the recurrent parent.

**[0008]** Three conventional approaches to interspecific hybridization for the purpose of alien transfer and incorporation of germplasm into stable diploid lines are (a)

diploid x diploid then doubling to produce the allopolyploid and backcrossing to produce the sesquidiploid, (b) autotetraploid x autotetraploid to produce the allopolyploid directly, and (c) autotetraploid x diploid to produce the sesquidiploid directly as described in Technical Bulletin 1586, U.S. Department of Agriculture, 1979.

**[0009]** Gene transfer mediated by *Agrobacterium tumefaciens* vectors has become routine in tobacco (DeBlock et al., 1987; Grierson, et al., 1990; Hilder, et al., 1990; Lindbo and Dougherty, 1992). Commercial use of genetic transformation in agriculture depends on the incorporation of foreign genes into high-yielding germplasm. Plant breeders may want to combine several foreign genes into a single elite germplasm source. In order to accomplish this goal, designer chromosome construction is desirable.

**[0010]** Designer, or artificial, chromosomes have been produced in yeast. In plants, the scaffold for designer chromosome construction can be found in a breeding line that possesses the full complement of chromosomes from its own species plus an addition chromosome from a related species, Campbell, et al., Construction of a designer chromosome in tobacco, *Theor Appl Genet* (1994) 87:837-842. Additional chromosomes are often meiotically stable as homozygotes, and because recombination between additional chromosomes and the rest of the plant genome is rare (Gerstel 1945) the integrity of a foreign-gene linkage package can be preserved.

**[0011]** The placement of this linkage package on an additional chromosome will also minimize disturbance to the rest of the plant genome. The genetic structure of high-yielding germ plasm can be disturbed either through insertional mutagenesis or the disturbance of beneficial linkage blocks. Insertional mutagenesis, caused by the integration of foreign genes into plant coding sequences, can occur frequently. Konez, et al. (1989) estimated that at least 30% of all T-DNA insertions occur in transcribed regions of the *Arabidopsis* and *Nicotiana* genomes. Beneficial linkage blocks are formed through intermating and recombination followed by selection.

**[0012]** The desired traits for a plant may not be present in the germplasm of the species of interest. In such a case, traditional breeding within the species may not give

acceptable results. Introducing a desired trait from one species into a related one by interspecific hybridization is followed by introgression into the recurrent parent.

**[0013]** Methods for expressing genes in plants has been described in U. S. Patent Numbers 5,316,931; 5,589,367; 5,811,653 and 5,866,785, all of which are incorporated herein. Certain species have either an acceptable biomass or an acceptable expression of a sequence inserted into a plant viral vector, but no current species have both high levels of biomass and high levels of viral vector performance. There is a need for a method to develop in one species both the characteristics of increased biomass and improved viral vector performance.

### SUMMARY OF THE INVENTION

**[0014]** The present invention is directed to a method of crossing two different species of *Nicotiana* to produce an interspecific hybrid which results in an increased biomass and improved host susceptibility to a viral vector.

**[0015]** In accordance with this invention, interspecific *Nicotiana* hybrids have been produced that are characterized with having a biomass of greater than that usually obtained from tobacco with CP-fusion constructs and a virion yield of greater than 1.0 mg. virion/gram fresh weight.

**[0016]** In one embodiment of the invention, there is provided a novel *Nicotiana* interspecific hybrid. This invention thus relates to the seeds of *Nicotiana* interspecific hybrid, to the plants of *Nicotiana* interspecific hybrid and to methods for producing a *Nicotiana* interspecific hybrid plant produced by crossing the *Nicotiana* interspecific hybrid with itself or another *Nicotiana* interspecific hybrid line.

**[0017]** Thus, any such methods using the *Nicotiana* interspecific hybrid are part of this invention: selfing, backcrosses, hybrid production, crosses to populations, and the like. All plants produced using a *Nicotiana* interspecific hybrid as a parent are within the scope of this invention. Advantageously, the *Nicotiana* interspecific hybrid could be used in crosses with other, different, *Nicotiana* interspecific hybrids to produce first generation ( $F_1$ ) *Nicotiana* interspecific hybrid seeds and plants with superior characteristics.

[0018] In another aspect, the present invention provides for single gene converted plants of *Nicotiana* interspecific hybrid. The single transferred gene may preferably be a dominant or recessive allele. Preferably, the single transferred gene will confer such traits as herbicide resistance, insect resistance, resistance for bacterial, fungal, or viral disease, male fertility, male sterility, enhanced nutritional quality, and industrial usage. The single gene may be a naturally occurring *Nicotiana* gene or a transgene introduced through genetic engineering techniques.

[0019] In another aspect, the present invention provides regenerable cells for use in tissue culture of a *Nicotiana* interspecific hybrid. The tissue culture will preferably be capable of regenerating plants having the physiological and morphological characteristics of the foregoing *Nicotiana* interspecific hybrid, and of regenerating plants having substantially the same genotype as the foregoing *Nicotiana* interspecific hybrid. Preferably, the regenerable cells in such tissue cultures will be embryos, protoplasts, meristematic cells, callus, pollen, leaves, anthers, roots, root tips, flowers, seeds, pods or stems. Still further, the present invention provides *Nicotiana* plants regenerated from the tissue cultures of the invention.

#### DETAILED DESCRIPTION OF THE INVENTION

[0020] To provide an understanding of several of the terms used in the specification and claims, the following definitions are provided:

[0021] Geneware™ viral vector - As used herein, Geneware™ viral vector refers to a system for expressing genes in plants, such as described in U. S. Patent Nos. 5,316,931; 5,589,367; 5,811,653 and 5,866,785, all incorporated herein by reference.

[0022] Biomass - As used herein, the term "biomass" means the total fresh weight in grams of plant tissue which is harvested above the ground. The terms "fresh weight" and "biomass" are used interchangeably herein.

[0023] Virion yield - As used herein the term "virion yield" means the total yield of the virus measured in mg of virion per gram fresh weight.

[0024] Host - As used herein, a "host" means a cell, tissue or organism capable of supporting replication of a vector or plant viral nucleic acid and which is capable of being infected by a virus containing the viral vector or plant viral nucleic acid.

**[0025]**      Infection - As used herein, an "infection" means the ability of a virus to transfer its nucleic acid to a host or introduce viral nucleic acid into a host, wherein the viral nucleic acid is replicated, viral proteins are synthesized, and new viral particles assembled. In this context, the terms "transmissible" and "infective" are used interchangeably herein.

**[0026]**      Phenotypic Trait - As used herein, "phenotypic trait" is an observable property resulting from the expression of a gene or genes.

**[0027]**      Plant Tissue - As used herein, "plant tissue" is any tissue of a plant in planting or in culture. This term is intended to include a whole plant, plant cell, plant organ, protoplast, cell culture, or any group of plant cells organized into a structural and functional unit.

**[0028]**      Recombinant Plant Viral Nucleic Acid - As used herein, "recombinant plant viral nucleic acid" means it has been modified to contain nonnative nucleic acid sequences.

**[0029]**      Recombinant Plant Virus - As used herein, "recombinant plant virus" is a plant virus containing the recombinant plant viral nucleic acid.

**[0030]**      Vector - As used herein, "vector" is a self-replicating nucleic acid molecule which transfers a nucleic acid segment between cells.

**[0031]**      Virus - As used herein, "virus" is an infectious agent composed of a nucleic acid encapsidated in protein. A virus may be a mono-, di-, tri- or multi-partite virus.

**[0032]**      Backcrossing - As used herein, "backcrossing" is a process in which a breeder repeatedly crosses hybrid progeny back to one of the parents, for example, a first generation hybrid  $F_1$  with one of the parental genotypes of the  $F_1$  hybrid.

**[0033]**      Essentially all the physiological and morphological characteristics - As used herein, a plant having essentially all the physiological and morphological characteristics means a plant having the physiological and morphological characteristics, except for the characteristics derived from the converted gene.

**[0034]**      Single Gene Converted (Conversion) - As used herein, "single gene converted (conversion)" plant refers to plants which are developed wherein essentially

all of the desired morphological and physiological characteristics of a variety are recovered in addition to the single gene transferred into the variety via the backcrossing technique or via genetic engineering.

**[0035]**     Trichomes - As used herein, "trichomes" means the glandular hairs on the surface of the plant stem, leaf, calyx, corolla or other plant tissue surfaces.

**[0036]**     Leaf lamina - As used herein, "leaf lamina" means the leaf surface devoid of vascular tissue of primary, secondary or ternary nature.

**[0037]**     GFP-GJ - As used herein, "GFP-GJ" means a crude plant extract containing green fluorescent protein.

**[0038]**     GFP-pH5, Δ - As used herein, "GFP-pH5, Δ" means a crude plant extract that was treated with heat and a pH of 5 that contains green fluorescent protein that is enriched for its presence.

**[0039]**     CP-GJ - As used herein, "CP-GJ" means a crude plant extract containing viral coat protein.

**[0040]**     CP-pH5, Δ - As used herein, "CP-pH5, Δ" means a crude plant extract that was treated with heat and a pH of 5 that contains coat protein that is enriched for its presence.

**[0041]**     Host plant species vary in their ability to support expression of a sequence inserted into a plant viral vector. For example, some species (such as *Nicotiana benthamiana*) support expression from a dual-subgenomic promoter tobamoviral vector to a high specific activity, but have relatively low biomass. Other species (such as *N. tabacum*) have high biomass and/or other desirable properties for growth in the field, but have a relatively low specific activity of the expressed protein. Using the method of the present invention, after chromosome doubling to restore fertility, the primary hybrid may have suitable properties, or it may be desirable to backcross toward either parent selecting or screening at each generation for the desired properties of the non-recurrent parent (e.g., introgressing the superior biomass of *N. tabacum* into *N. benthamiana*, or introgressing the superior viral vector performance of *N. benthamiana* into *N. tabacum*). A viral vector expressing the green fluorescent protein (GFP) is one example of a useful tool for screening the level of systemic expression in candidate hybrid plants.

Geneware™ viral vector refers to a system for expressing genes in plants such as described in U. S. Patent Nos. 5,316,931; 5,589,367; 5,811,653 and 5,866,785, all incorporated herein by reference.

### EXAMPLES

**[0042]** The following examples are provided to further illustrate the present invention and are not intended to limit the invention beyond the limitations set forth in the appended claims.

#### Example 1

##### Procedures for Making Interspecific Crosses of *Nicotiana*

**[0043]** When making interspecific crosses it is important to have: 1) healthy plants, 2) careful emasculation, 3) freedom from contamination by foreign pollen, 4) a reasonable level of cross fertility among the parent plants, and 5) recording of the crosses. A pair of fine curved-point forceps is used to slit the corolla and emasculate the blossom. In the case of *N. tabacum*, and many other species, the anthers are removed just before the corolla unfolds. Several exceptions apply to this general rule. In the case of *N. repanda*, the anthers are adnate to the stigma, the anther walls usually break down, and the contents of the anther adhere to the stigma before the flower matures. Anthers in the later-formed flowers of *N. nudicaulis* may dehisce before the corolla unfolds and self-pollination will occur. The flowers of these species are emasculated several days before cross-pollinations are made.

**[0044]** Ordinarily, pollinations are made immediately after emasculation, up to 18 to 24 hours after emasculation before applying pollen. The glistening and sticky surface of the stigma is touched with a newly dehisced anther, or pollen may be applied with the aid of a small artist's brush (size 6/128 to 10/128). Brush pollination aids seed set in some self-fertile species, and sib-pollinations are required to produce seed of self-sterile species. The brush used to transfer pollen is inserted between siblings, handle down, in a pot. Inexpensive brushes with plastic handles are preferable to those with wooden handles as microorganismal activity will cause the latter to rot. If pollen of a desired parent is unavailable from living plants, use pollen that was collected at an



earlier time and stored frozen in gelatin capsules. Gwynn has found that frozen pollen from *N. tabacum* may remain viable for as long as 7 years.

**[0045]** If a parental plant is suspected of harboring a mechanically transmitted virus, the instruments are sterilized in 70 percent ethanol. After a pollination has been made, a length of drinking straw, slightly longer than the pistil, is placed over the stigma and crimped by folding to prevent contamination by self- or foreign-pollen. Use of straws is probably unnecessary if plants are well spaced on the greenhouse bench, mature blossoms are removed, and neither insects nor excessive air movements present a problem. Furthermore, under most circumstances, a stigma thoroughly covered with the intended pollen offers sufficient protection against contamination. Usually, fertilization is affected within 1 to 2 days. Mark individual crosses with a small, stringed key tag. Moisten the string before looping it around the pedicel otherwise the flower may be severed when the string is tightened. Whenever a number of identical crosses are planned, time can be saved by marking the tags in advance of pollination.

**[0046]** Generally, seed of most *Nicotiana* spp. will mature about 28 days after pollination. Seedlings of *N. tabacum* may be obtained earlier by scraping the developing seeds from the carpels about 18 days after pollination and treating them in 0.5 percent sodium hypochlorite (Clorox and water, 1:9) for about 10 minutes. One or two capsules of *N. tabacum* will contain enough seed to maintain a breeding line, but the seed yield per capsule of other species may be considerably smaller.

**[0047]** Some *Nicotiana* spp. are essentially day neutral whereas others require special temperature and day-length conditions to initiate flowering. Therefore, when planning specific hybridizations, consider the day-length requirements of the species involved, unless you can freeze pollen for use at a later time. Certain crosses may require a multitude of individual pollinations with the hope of obtaining a single seed among many mature capsules, for example, the cross diploid *N. rustica* x diploid *N. tabacum*. Difficult hybridizations may yield a paucity of shrunken seed of a smaller than average size. Although most seeds of this kind are often inviable, some of them may be induced to germinate if they are sown on soil in covered plastic or glass preparation dishes. when seeds are limited in number, do not treat the seedlings with colchicine.

Selecting the most vigorous plants is better for asexual propagation. Later, treat the growing points of the clonal material with 0.5 percent aqueous colchicine three times a day for 3 days to induce chromosome doubling.

**[0048]** Premature blossom drop may be avoided by the use of 0.5 percent indoleacetic acid in lanolin. Apply it by dipping a needle into the preparation and scratching the pedicel at its point of attachment. This treatment will cause the blossom to remain on the plant regardless of embryo development and, therefore, offers no guarantee of a successful hybridization. On the other hand, syngamy serves to prevent abscission.

### Example 2

#### Selection of Parental Lines for *N. excelsior*/*N. benthamiana* Hybrids

**[0049]** Seven accessions of *N. excelsior* were collected. All of the accessions had similar characteristics, but only TW46 and TW47 were determined to be good hosts for GFP Geneware™ viral vector. TW46 was identified as a poor host for certain insert genes in Geneware™ viral vector. Since systemic necrosis was observed in TW46, TW47 was selected. A group of over 100 *N. excelsior* TW47 were inoculated to give a statistically valid survey of the performance of GFP Geneware™ viral vector on *N. excelsior*. Almost all of the plants gave good-to-excellent systemic GFP expression. A virus preparation from 200 grams of infected whole plants gave a final yield of 2 mg virion/g fresh weight. Unexpectedly this was 5- to 10-fold higher than the virion yield typically obtained from tobacco with CP-fusion constructs. The higher virion yield for *N. excelsior* when extended to the field, the yield per acre would be similar to tobacco, but the amount of tissue (waste) in the bioprocessing stream is greatly reduced. Typical results for *N. benthamiana* are 0.9 mg virion/g fresh weight.

**[0050]** Seven accessions of *N. benthamiana* were screened for biomass and as a host for systemic expression from dual-subgenomic Geneware™ viral vector. One of the best performers was TW16.

### Example 3

**[0051]** A primary interspecific hybrid between *N. excelsior* TW47 and *N. benthamiana* TW16 gave very good systemic GFP expression with GFP Geneware™

**[0052]** Several regenerants set seed indicating chromosome-doubling and restored fertility. The working name for the hybrid species is *N. excelsiana*. The seed was germinated, and the resulting plants were evaluated for biomass and systemic Geneware™ viral vector expression. Systemic GFP expression was near the level of *N. benthamiana*. Biomass and growth habit were similar to *N. excelsior*.

**[0054]** The primary chromosome-doubled hybrid between *N. excelsior* and *N. benthamiana* (*N. excelsiana*) appears to be a stable true-breeding line. It has good biomass and excellent Geneware™ viral vector properties, and did well in a test in Kentucky.

**[0055]** In Table 1 listed below the phenotypic observations for several traits are shown for two lines each of the parents *Nicotiana benthamiana* TW16; *Nicotiana excelsior* TW47 and the hybrid *Nicotiana excelsiana* 7.b.

TABLE 1

Phenotypic Observations Planted 3/10/99; Data Collected 4/22/99						
Traits	<i>Nicotiana benthamiana</i>		<i>Nicotiana excelsior</i>		<i>Nicotiana excelsiana</i>	
	TW16 #1	TW16 #2	TW47 #1	TW47 #2	7.b #1	7.b #2
First flower open	4/23/99	4/22/99	Budding	Budding	Budding	Budding
Height (to top of stem)	20 cm	26 cm	27 cm	20 cm	34 cm	26 cm
Largest leaf (Length)	18 cm	19.5 cm	17-20 cm	19.5 cm	21 cm	19 cm
Largest leaf (Width)	13 cm	13.5 cm	10-12 cm	10 cm	14 cm	13 cm
Leaf shape	Round	Round	Oblong	Oblong	Oval	Oval
Leaf angle	90°	90°	90°	90°	90°	90°
Leaf color	Med. green	Med. green	Dark green	Dark green	Med. green	Med. green
Trichomes	Mod. #, small	Mod. #, small	Few, large	Few, large	Few, large	Few, large
Petiole leaf	No	No	Yes, 70-80mm	Yes, 40mm	Yes, 20mm	Yes, 14mm
Petiole length	5 cm	6 cm	Continuous w/ leaf lamina	Continuous w/ leaf lamina	6 cm	5 cm
Leaf lamina	13 x 13 cm	13.5 x 13.5 cm	20 x 12 cm	19.5 x 10 cm	15 x 14 cm	14 x 13 cm
Internode length	10 mm	15 mm	15 mm	15 mm	22 mm	21 mm
Stalk diameter	7 mm	7 mm	7 mm	8 mm	7 mm	7 mm
Branching	Yes @ every leaf node	Yes @ every leaf node	Yes, barely from 5 leaf nodes	Not yet	Yes, 2 nodes starting to extend	Yes, 2 nodes starting to extend
Flower form	Nb	Nb				
Flower color	White	White				
Flower head habit	1 flower/leaf node on main stem	1 flower/leaf node on main stem				

TABLE 2

Phenotypic Observations Planted 3/10/99; Data Collected 4/26/99						
Traits	<i>Nicotiana benthamiana</i>		<i>Nicotiana excelsior</i>		<i>Nicotiana excelsiana</i>	
	TW16 #1	TW16 #2	TW47 #1	TW47 #2	7.b #1	7.b #2
First flower open	Yes, secondary meristems starting bud	Yes, secondary meristems starting bud	4/25/99 Buds on topmost branches	4/25/99 Buds on topmost branches	4/25/99 Buds on topmost branches	Budding; Estd opening 4/28/99
Height (to top of stem)	24 cm	31 cm	40 cm	31 cm	49 cm	40 cm
Largest leaf (Length)	21 cm	21 cm	21 cm	21 cm	23 cm	22 cm
Largest leaf (Width)	15 cm	15 cm	14 cm	11.5 cm	15.5 cm	13.7 cm
Leaf shape	Round	Round	Oblong	Oblong	Oval	Oval
Leaf angle	90°	90°	90°	<90°	90°	90°
Leaf color	Med. green	Med. green	Dark-med. green	Med. green	Med. green	Med. green
Trichomes	Small, med. #	Small, med. #	Few, large	Few, large	Few, large	Few, large
Petiole leaf	No	No	Yes, 9.5 cm	Yes, 4 cm	Yes, 20 mm	Yes, 18 mm
Petiole length	5 cm	6 cm	N/A	N/A	7 cm	5.5 cm
Leaf lamina	16 x 15 cm	15 x 15 cm	21 x 14 cm	21 x 11.5 cm	16 x 15.5 cm	16.5 x 13.7 cm
Internode length	1.6 cm (10 nodes)	2 cm (10 nodes)	3.2 cm (5 nodes)	2.9 cm (5 nodes)	3 cm (8 nodes)	2.4 cm (10 nodes)
Stalk diameter	7 mm	8 mm	8 mm	7 mm	8 mm	8 mm
Branching	Yes, almost all leaf nodes	Yes, almost all leaf nodes	Yes, 5 are extending	Yes, 5 are extending	Yes, 9 are extending	Yes, 6 are extending
Flower form	Nb	Nb	Large	Large	Medium	N/A
Flower color	White	White	White	White	White	N/A
Flower head habit	1/node from main stalk	1/node from main stalk	1/node from main stalk	1/node from main stalk	1/node from main stalk	N/A

TABLE 3

Plant	GFP-GJ	GFP-pH5, Δ	CP-GJ	CP-pH5, Δ
<i>N. benthamiana</i> lab strain	0.5 mg/g FW	0.6 mg/g FW	1.1 mg/g FW	1.6 mg/g FW
<i>N. tabacum</i> MD609	ND	ND	ND	ND
BB/TT hybrid	0.2	0.1	0.2	0.1
<i>N. benthamiana</i> TW16	1.0	0.8	1.4	1.2
<i>N. excelsior</i> TW47	1.2	0.8	2.6	2.4
<i>N. excelsiana</i>	0.8	0.6	1.9	2.2
<i>N. occidentalis</i> TW91	0.4	0.6	1.5	1.7
<i>N. umbratica</i> TW144	0.9	0.9	1.0	1.0
TW91 x TW144	0.5	0.5	0.8	0.9
TW144 x TW91	0.7	0.6	1.3	1.3

#### Example 4

##### Backcross Progeny from Interspecific Hybrid TW47 x TW16

[0056] Backcrosses to each parental type were performed in order to evaluate the characteristics of the sesquidiploids. The backcross to *N. excelsior* was the only one to germinate. They looked much like *N. excelsior* and gave a similar GFP expression pattern to *N. excelsior*. An additional backcross was performed to generate the first breakdown generation, and the resulting plants are currently being evaluated.

#### Example 5

[0057] In-field screening of various *Nicotiana* species was continued in 1999. The purpose of this study was to evaluate biomass differences between species, to evaluate growth characteristics differences between species, and to evaluate insect and disease pressure differences between species.

[0058] Nine different *Nicotiana* species were seeded in the greenhouse on April 14. Each species was seeded into three 72-cell trays and placed in a specially constructed float bed. Burley Gold growth media was used for the transplants. Trays were filled manually and dibbled using the Golf Ball Dibbler. Raw seed was used for eight of the species and pelleted *Benthamiana* seed was used for the lab strain *Benthamiana* transplants. The GROmore hand seeder was used to sow the raw seed and the pelleted seed was sown by hand. Each tray was labeled with a tray marker to identify each species. The pelleted lab strain *Benthamiana* trays were watered overhead for three days post-seedling to dissolve the clay pellet coating and facilitate cotyledon emergence. Table 4 outlines the species transplanted in this study.

Table 4

<b><i>Nicotiana</i> Species Transplanted</b>	
<b><i>Nicotiana</i> Species</b>	<b>Species Identifier</b>
TW 17 - <i>Nicotiana Benthamiana</i>	Harvested 12/8/97
TW 16 - <i>Nicotiana Benthamiana</i>	Harvested 11/26/97
SCR1 - <i>Nicotiana Benthamiana</i>	Harvested 11/6/98
L3663 - <i>Nicotiana Benthamiana</i>	Harvested 4/4/98
<i>Nicotiana Excelsiana</i>	Lot #7.b, Harvested 5/22/98
<i>Nicotiana Occidentalis Obliqua</i>	Unknown SCRI Stock
TW 47 - <i>Nicotiana Excelsior</i>	Harvested 3/31/98
TW 91 - <i>Nicotiana Occidentalis</i>	Harvested 12/16/97
Lab Strain - <i>Nicotiana Benthamiana</i>	Lot #28b 29b RC

**[0059]** Transplants were treated, through the float water, with the fungicide Ridomil. An insecticide, Orthene, and a foliar applied fungicide, Dithane were also applied to the transplants. At 21 DPT, fertilizer was applied to the float water. A fertility level of 7 on the DiSST-4 meter was maintained throughout transplant production.

**[0060]** Transplants were transported to the ARP for transplanting on May 20, 37 DPS. Plants were transplanted on raised, black plastic covered beds, using a mechanical transplanter at a density of four plants per linear foot of bed. The plants were transplanted in two rows on the bed with an offset plant configuration. The transplant water contained ½ gallon of 7-14-7 Starter Fertilizer and ½ pound of Orthene per 150 gallons of water. Each species was replicated three times with each replication consisting of ten feet of bed.

**[0061]** At transplanting, several differences were apparent in transplant quality. TW 16 produced a slightly leggy transplant that wilted soon after transplanting. TW 17 transplants were smaller than the other *Benthamiana* transplants. However, TW 17 transplants had a thicker stem and appeared much hardier than the other *Benthamiana* transplants. The Lab Strain *Benthamiana* plants were much smaller and less



**[0062]** All plot replications were treated every 7-10 days with Orthene and Dithane at a rate of 1 teaspoon/gallon. Soil moisture was monitored using gypsum blocks that were buried about 4-5 inches in the beds. No irrigation was necessary for this plot due to the timely rainfall.

**[0063]** On June 29, 43 DPT, four plants from each replication were sampled for fresh weight determination. Table 5 illustrates the results for each replication.

Tabl 5

Fresh Weights for each Replication		
Species	Replication No.	Weight/Plant (g/plant)
TW 16 <i>Benthamiana</i>	1	411.6
TW 16 <i>Benthamiana</i>	2	235.7
TW 16 <i>Benthamiana</i>	3	280.1
TW 17 <i>Benthamiana</i>	1	426.8
TW 17 <i>Benthamiana</i>	2	352.6
TW 17 <i>Benthamiana</i>	3	Dead
SCRI <i>Benthamiana</i>	1	326.3
SCRI <i>Benthamiana</i>	2	236.6
SCRI <i>Benthamiana</i>	3	Dead
L3663 <i>Benthamiana</i>	1	335.6
L3663 <i>Benthamiana</i>	2	292.1
L3663 <i>Benthamiana</i>	3	Dead
TW 47 <i>Excelsior</i>	1	616.5
TW 47 <i>Excelsior</i>	2	450.6
TW 47 <i>Excelsior</i>	3	517.8
TW 91 <i>Occidentalis</i>	1	528.8
TW 91 <i>Occidentalis</i>	2	399.8
TW 91 <i>Occidentalis</i>	3	449.2
<i>Occidentalis Obliqua</i>	1	141.9
<i>Occidentalis Obliqua</i>	2	620.6
<i>Occidentalis Obliqua</i>	3	809.7
<i>Excelsiana</i>	1	333.0
<i>Excelsiana</i>	2	512.4
<i>Excelsiana</i>	3	580.8
Lab Strain <i>Benthamiana</i>	1	90.5
Lab Strain <i>Benthamiana</i>	2	200.5
Lab Strain <i>Benthamiana</i>	3	N/A

**[0064]** Several replications died due to root rot induced by excess water surrounding the raised bed. This excess water came about due to excessive rainfall and poor plot drainage.

**[0065]** Table 6 summarizes the replication plant weights. The following Table 6 is the average of all replications for each species.

**Table 6**

<b>Replication Averages by Species</b>	
<b>Species</b>	<b>Average Plant Weight (g/plant)</b>
TW 16 <i>Benthamiana</i>	305.8
TW 17 <i>Benthamiana</i>	389.7
SCRI <i>Benthamiana</i>	281.5
L3663 <i>Benthamiana</i>	313.9
TW 47 <i>Excelsior</i>	528.3
TW 91 <i>Occidentalis</i>	459.3
<i>Occidentalis Obliqua</i>	524.1
<i>Excelsiana</i>	475.4
Lab Strain <i>Benthamiana</i>	145.5

**[0066]** TW 17 performed well in comparison to the other *Benthamiana* in the plot. TW 17 was a very vigorous plant with excellent hardiness and high biomass levels. More work should be done to determine the feasibility of this plant as a viral vector host. Also, *Excelsiana* stood out as a superior plant in this plot. This plant offered high biomass levels with significant leaf size and weight.

**[0067]** This invention is also directed to methods for producing a *Nicotiana* plant by crossing a first parent *Nicotiana* plant with a second parent *Nicotiana* plant, wherein the first or second *Nicotiana* plant is the *Nicotiana* plant from the line *Nicotiana* Interspecific Hybrid. Further, both first and second parent *Nicotiana* plants may be from the cultivar TW47 x TW16. Therefore, any methods using this cultivar are part of this

invention: selfing, backcrosses, hybrid breeding, and crosses to populations. Any plants produced using cultivar as a parent are within the scope of this invention.

**[0068]** As used herein, the term "plant" includes plant cells, plant protoplasts, plant cells of tissue culture from which *Nicotiana* plants can be regenerated, plant calli, plant clumps, and plant cells that are intact in plants or parts of plants, such as pollen, flowers, embryos, ovules, seeds, pods, leaves, stems, roots, anthers and the like. Thus, another aspect of this invention is to provide for cells which upon growth and differentiation produce a cultivar having essentially all of the physiological and morphological characteristics of *Nicotiana* Interspecific Hybrid.

**[0069]** Culture for expressing desired structural genes and cultured cells are known in the art. Also as known in the art, *Nicotiana* are transformable and regenerable such that whole plants containing and expressing desired genes under regulatory control may be obtained. General descriptions of plant expression vectors and reporter genes and transformation protocols can be found in Gruber, et al., "Vectors for Plant Transformation, in Methods in Plant Molecular Biology & Biotechnology" in Glich, et al., (Eds. pp. 89-119, CRC Press, 1993). Moreover GUS expression vectors and GUS gene cassettes are available from Clone Tech Laboratories, Inc., Palo Alto, California while luciferase expression vectors and luciferase gene cassettes are available from Pro Mega Corp. (Madison, Wisconsin). General methods of culturing plant tissues are provided for example by Maki, et al., "Procedures for Introducing Foreign DNA into Plants" in Methods in Plant Molecular Biology & Biotechnology, Glich, et al., (Eds. pp. 67-88 CRC Press, 1993); and by Phillips, et al., "Cell-Tissue Culture and In-Vitro Manipulation" in Corn & Corn Improvement, 3rd Edition; Sprague, et al., (Eds. pp. 345-387) American Society of Agronomy Inc., 1988. Methods of introducing expression vectors into plant tissue include the direct infection or co-cultivation of plant cells with *Agrobacterium tumefaciens*, Horsch et al., Science, 227:1229 (1985). Descriptions of *Agrobacterium* vectors systems and methods for *Agrobacterium*-mediated gene transfer provided by Gruber, et al., supra.

**[0070]** Useful methods include but are not limited to expression vectors introduced into plant tissues using a direct gene transfer method such as microprojectile-mediated delivery, DNA injection, electroporation and the like. More preferably expression vectors are introduced into plant tissues using the microprojectile media delivery with the biolistic device *Agrobacterium*-mediated transformation. Transformant plants obtained with the protoplasm of the invention are intended to be within the scope of this invention.

**[0071]** The present invention contemplates a *Nicotiana* plant regenerated from a tissue culture of a variety or hybrid plant of the present invention. As is well known in the art, tissue culture of *Nicotiana* can be used for the in vitro regeneration of a *Nicotiana* plant. Tissue culture of various tissues of *Nicotiana* and regeneration of plants therefrom is well known and widely published. For example, reference may be had to Weissbach and Weissbach (eds.), Meth. Enzymol. 118 (1986); Zaitlin et al. (eds), Biotechnology in Plant Science (1985).

**[0072]** When the term *Nicotiana* plant is used in the context of the present invention, this also includes any single gene conversions of that variety. The term single gene converted plant as used herein refers to those *Nicotiana* plants which are developed by a plant breeding technique called backcrossing wherein essentially all of the desired morphological and physiological characteristics of a variety are recovered in addition to the single gene transferred into the variety via the backcrossing technique. Backcrossing methods can be used with the present invention to improve or introduce a characteristic into the variety. The term backcrossing as used herein refers to the repeated crossing of a hybrid progeny back to the recurrent parent. The parental *Nicotiana* plant which contributes the gene for the desired characteristic is termed the nonrecurrent or donor parent. This terminology refers to the fact that the nonrecurrent parent is used one time in the backcross protocol and therefore does not recur. The parental *Nicotiana* plant to which the gene or genes from the nonrecurrent parent are transferred is known as the recurrent parent as it is used for several rounds in the backcrossing protocol (Poehlman & Sleper, 1994; Fehr, 1987). In a typical backcross protocol, the original variety of interest (recurrent parent) is crossed to a second variety

(nonrecurrent parent) that carries the single gene of interest to be transferred. The resulting progeny from this cross are then crossed again to the recurrent parent and the process is repeated until a *Nicotiana* plant is obtained wherein essentially all of the desired morphological and physiological characteristics of the recurrent parent are recovered in the converted plant, in addition to the single transferred gene from the nonrecurrent parent.

**[0073]** The selection of a suitable recurrent parent is an important step for a successful backcrossing procedure. The goal of a backcross protocol is to alter or substitute a single trait or characteristic in the original variety. To accomplish this, a single gene of the recurrent variety is modified or substituted with the desired gene from the nonrecurrent parent, while retaining essentially all of the rest of the desired genetic, and therefore the desired physiological and morphological, constitution of the original variety. The choice of the particular nonrecurrent parent will depend on the purpose of the backcross, one of the major purposes is to add some commercially desirable, agronomically important trait to the plant. The exact backcrossing protocol will depend on the characteristic or trait being altered to determine an appropriate testing protocol. Although backcrossing methods are simplified when the characteristic being transferred is a dominant allele, a recessive allele may also be transferred. In this instance it may be necessary to introduce a test of the progeny to determine if the desired characteristic has been successfully transferred.

**[0074]** Many single gene traits have been identified that are not regularly selected for in the development of a new variety but that can be improved by backcrossing techniques. Single gene traits may or may not be transgenic, examples of these traits include but are not limited to, male sterility, herbicide resistance, resistance for bacterial, fungal, or viral disease, insect resistance, male fertility, enhanced nutritional quality, industrial usage, yield stability and yield enhancement. These genes are generally inherited through the nucleus.

DEPOSIT INFORMATION

**[0075]** A deposit of the *Nicotiana* seed of this invention has been made with the American Type Culture Collection, Manassas, Virginia. The deposit was made on July 12, 1999 and has ATCC Accession No. PTA-323.

**[0076]** Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity and understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the invention, as limited only by the scope of the appended claims.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000